



## ORIGINAL ARTICLE


Nutrition Education and Dietetics Infant, Child, and Adolescent Nutrition

# Prevalence of anemia, hematocrit variations, and micronutrient supplementation practices among pregnant women attending antenatal clinics in Ibadan, Southwest Nigeria

Rukiyat Abdus-salam<sup>1</sup> , Olaolu Oni<sup>1</sup> , Ayodeji Adeyanju<sup>2</sup> , Oluwabunmi Adeyeye<sup>2</sup> , Ajani Victoria<sup>2</sup>, Taofeeq Oluwatosin<sup>2</sup>, Mercy Adebayo-Tayo<sup>2</sup> , Obiageli Medebem<sup>2</sup> , Esosa Tongo<sup>2</sup> , Adenike Ajagbe<sup>2</sup> , Amarachi Nwaigwe<sup>2</sup> , Taiwo Akinosi<sup>2</sup> , Olatunbosun Oladayo<sup>2</sup>, Medical Students Group B 2019<sup>2</sup>

<sup>1</sup> Department of Obstetrics and Gynecology, College of Medicine, University College Hospital/University of Ibadan, Ibadan, Nigeria

<sup>2</sup> Department of Obstetrics and Gynecology, University of Ibadan, Ibadan, Nigeria



## ABSTRACT

**Background:** Anemia is a prevalent complication of pregnancy that poses significant a risk to both maternal and fetal health, potentially leading to adverse pregnancy outcomes. According to the World Health Organization (WHO), anemia is diagnosed when hemoglobin (Hb) levels are below 11g/dL, with nutritional deficiencies being the most common underlying cause.

**Aim:** This study aimed to evaluate the prevalence of anemia, the intake of common micronutrient supplements, and other factors influencing changes in hematocrit levels among pregnant women.

**Patients and Methods:** A cross-sectional study was conducted in antenatal clinics in Ibadan from January to February 2023. Data were collected from 309 pregnant women in their third trimester using a pre-tested, interviewer-administered questionnaire. The questionnaire captured socio-demographic details, obstetric characteristics, packed-cell volume, and the intake of micronutrient supplements. Descriptive statistics were performed, and paired sample t-test along with linear regression analyses were used to determine associations. Statistical significance was set at  $p < 0.05$ .

**Results:** The mean age of the participants was  $30.49 \pm 5.03$  years, with the majority having tertiary education (77%) and identifying as Christians (57%). At the time of booking, 42.9% of participants were anemic, with a mean packed cell volume (PCV) of 33.46% (SD = 3.83). A significant reduction in anemia prevalence was observed during the interview, with 26.8% of participants classified as anemic (Mean PCV = 34.03, SD=2.90) [ $t(155) = 2.089, p = 0.038$ ]. While 73.5% of participants reported adequate folic acid intake, only 51.1% consumed sufficient ferrous tablets. Fewer than 30% of participants reported adequate intakes of calcium, vitamin A and other micronutrients. Changes in hematocrit levels were not significantly associated with micronutrient intake but were significantly influenced by participants' age group ( $p = 0.029$ ), level of education ( $p = 0.041$ ), and religion ( $p = 0.007$ ).

**Conclusion:** The study observed a significant reduction anemia prevalence during the third trimester among participants, despite suboptimal micronutrient supplementation. This suggests the possible contribution of alternative sources of micronutrients, likely dietary, with age and educational attainment emerging as significant influencing factors.

**Keywords:** Anemia, hematocrit changes, micronutrients supplementation, pregnancy, women.

## ARTICLE INFORMATION

**Corresponding author:** Olaolu Oni.

**E-mail:** olaoluoni43@yahoo.com

**Tel.** +234 (8062087527)

**Received:** July 19, 2024

**Revised:** October 16, 2024

**Accepted:** January 11, 2025

**Published:** January 11, 2025

**Article edited by:**

Prof. Khaled Méghit Boumédiène

**Article reviewed by:**

Dr. Asha Murshida Muthalib

Dr. Prosper Kujinga Chopera

Cite this article as: Abdus-salam R., Oni O., Adeyanju A., Adeyeye O., Victoria A., Oluwatosin T., Adebayo-Tayo M., Medebem O., Tongo E., Ajagbe A., Nwaigwe A., Taiwo A., Oladayo O., Medical Students Group B 2019 (2025). Anemia, hematocrit changes and micronutrients supplementation among antenatal attendees in Ibadan, southwest, Nigeria. *The North African Journal of Food and Nutrition Research*, 9(19): 1–12. <https://doi.org/10.51745/najfnr.9.19.1-12>

© 2025 The Author(s). This is an open-access article. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

## 1 Introduction

Anemia is defined as a reduction in the level of red blood cells or the oxygen-carrying protein hemoglobin (Hb) within these cells (World Health Organization [WHO], 2011). Pregnant women and children are particularly vulnerable to anemia due to increased physiological demand and nutritional deficiencies during pregnancy (Galloway, 2003). This

condition poses significant risk to both maternal and fetal health on a global scale (Agbozo *et al.*, 2020). The WHO defines anemia in pregnancy as a Hb level below 11g/dL. Hemoglobin levels vary across the trimesters due to physiological changes, such as increased plasma volume and hemodilution, which are more pronounced in the second trimester, as well as the heightened nutritional demands

imposed by fetal growth in the later stages of pregnancy (Chowdhury, 2015).

Globally, anemia affects approximately 41.8% of pregnancies, with a higher prevalence of 46% observed in Africa, where iron deficiency accounts for the majority of cases (WHO, 2020). The prevalence of anemia ranges from 15–19% in high-income countries to between 33% and 75% in low- and middle-income countries (Abdallah *et al.*, 2022; Cappellini & Motta, 2015).

In sub-Saharan Africa, the primary cause of anemia is nutritional deficiency (Chowdhury, 2015). Other contributing factors include hemoglobinopathies, severe infection during pregnancy, chronic medical conditions, and malaria (Kidanto *et al.*, 2009). Among nutritional deficiencies, iron deficiency anemia (IDA) is the most prevalent. The increased resorption and utilization of iron stores during pregnancy exacerbate the condition, particularly for women with inadequate iron reserves before conception. Women with depleted iron stores may struggle to meet the heightened iron requirements during pregnancy, necessitating supplementation of micronutrients to address nutritional deficiencies (Barrett *et al.*, 1994).

An imbalance in the intake of macro- and micro-nutrients can result in deficiencies, which are significant contributors to nutritional disorders, including anemia—especially folate and iron deficiency anemia. Surveys conducted in Ethiopia, Kenya, Nigeria, and South Africa have reported the prevalence of deficiencies in vitamin A, vitamin B12, iodine, zinc, and folate among pregnant women as 21–48%, 8–10%, 87%, 46–76% and 3–12% respectively (Frayne *et al.*, 2014; Harika *et al.*, 2017).

Micronutrient supplementation refers to the provision of additional nutrients, such as vitamins and minerals, to address dietary inadequacies. During pregnancy, the demand for micronutrients increases significantly to meet maternal and fetal requirements. However, women in low- and middle-income countries often fail to achieve adequate micronutrient intake (Frayne *et al.*, 2014; Harika *et al.*, 2017). Beyond the use of iron supplements to prevent anemia, reduce the risk of intrauterine growth restriction, and lower the likelihood of low birth weight, other essential micronutrients are recommended. These include folic acid to prevent neural tube defects, iodine to mitigate cretinism, zinc to reduce the risk of preterm birth, and vitamin D alongside folic acid to improve birth weight (Gernand *et al.*, 2016).

Women in low-income countries are particularly vulnerable to micronutrient deficiencies due to limited access to supplements, insufficient consumption of fortified foods, as well as the frequent occurrence of infectious diseases (Darnton-Hill, 2012). The WHO advocates for pregnant women to receive adequate nutrition and encourages the use

of micronutrient supplements, including iron, folic acid, vitamins A and B, and calcium, as required (WHO, 2020). Consistent use of these supplements enhances maternal micronutrient stores and ensures their availability for the developing fetus.

Despite these recommendations, compliance with micronutrient supplementation during pregnancy remains suboptimal. For instance, a survey conducted in Nigeria revealed that only 21% of women took iron tablets daily for 90 or more days during their last pregnancy (National Population Commission (NPC) [Nigeria] and ICF International, 2014). While numerous studies have examined the relationship between single micronutrient supplementation—especially iron—and hematocrit levels during pregnancy, and showed a paucity of research evaluating the effect of multiple micronutrient supplementation. Hence, it is important to evaluate the use of various micronutrients, compliance benefits and barriers to use among pregnant women. This study, therefore, assessed the prevalence of anemia, the intake of commonly recommended micronutrient supplements, and the factors associated with changes in hematocrit level during pregnancy.

## 2 Methods

This cross-sectional study was carried out in the antenatal clinics of University College Hospital (UCH) and Adeoyo Maternity Hospital, both located in Ibadan-North Local Government Area of Oyo State, South-west, Nigeria. The study period spanned January to February 2023.

Eligible participants included pregnant women in their third trimester who were receiving antenatal care at the selected study sites. Inclusion criteria required participants to provide informed consent and meet the study's predefined eligibility requirements. Pregnant women with hemoglobinopathies, chronic infections, antepartum hemorrhage, chronic medical conditions, multiple pregnancies, or a history of allergies to iron or other micronutrients were excluded.

The sample size was calculated using the Leslie Kish formula (Kish, 1965), with a prevalence of anemia of 25.6% from a previous study (Abdallah *et al.*, 2022),  $Z\alpha = 1.96$ , and an allowance for 10% attrition. The resulting required sample size was 293 participants; however, 309 participants were ultimately recruited to enhance the study's statistical power. All participants were thoroughly briefed on the study's objectives and procedures, and written informed consent was obtained before enrollment.

Data collection was conducted using a pre-tested, semi-structured, interviewer-administered questionnaire. The questionnaire gathered information on socio-demographic and obstetric characteristics, participants' packed cell volume (PCV) at their first antenatal clinic visit (booking) and at the

time of study enrolment, the pattern of intake of various micronutrient supplements, and other medications taken by the pregnant woman.

For pregnant women taking prenatal vitamins containing combined/multiple micronutrient supplements, such as “Pregncare<sup>®</sup>”, “Astyfer<sup>®</sup>”, “Obtron-6<sup>®</sup>”, etc., the specific micronutrient composition of these drugs was reviewed and recorded individually. This was achieved by examining and extracting information from the respective drug leaflets. The average elemental iron content in these hematinic supplements ranged from 17 mg to 60 mg, with the WHO, (2024) recommending a minimum daily requirement of 30 mg. The folic acid content ranged from 400 µg to 5000 µg, with a minimum daily requirement of 400 µg for non-anemic individuals (WHO, 2024).

The PCV assessment was conducted as a part of routine antenatal care. Capillary blood samples were obtained via finger prick, and centrifugation was performed using a micro-hematocrit centrifuge at 3000 rpm for five minutes. The PCV was then measured using a Hewkley micro-hematocrit reader. For this study, a PCV of less than 33% was categorized as indicative of anemia during pregnancy.

The intake pattern of specific micronutrient supplements, including iron, folic acid, and vitamin B, calcium, zinc, and vitamin A, supplements were categorized into two groups: *adequate intake* (optimal) and *adequate intake*. Adequate intake was defined as consuming the supplements at least four days per week, while inadequate intake was defined as consuming them three days or fewer per week. Dietary intake was assessed using the Spanish Food Frequency Questionnaire (FFQ), that was validated and adapted for use in the local context (Ruiz-Cabello et al., 2017).

The data collected were entered and analyzed using SPSS version 25. Descriptive statistics were performed, and the results were presented in tables displaying frequency counts, percentages and means. To examine the association between participants’ demographic and obstetrics characteristics and changes in PCV levels, linear regression for repeated measures was applied. The study center was treated as a control factor, while fixed factors included participants’ age, marital status, level of education, income, religion, ethnicity, gravidity, parity, and trimester.

The level of significance was set at  $p < 0.05$ , with a 95% confidence interval. However, given the multiple tests performed on the outcome variables, a family-wise error rate was calculated to be 14.3%. Consequently, the initial  $p$ -value ( $\alpha$ ) of 0.05 was adjusted to 0.017 using the Bonferroni correction to account for this error rate.

Ethical approval for the study was obtained from the Joint University of Ibadan and University College Hospital Ethics Review Committee (UI/UCH Ethics Committee) under approval number UI/EC/23/0040.

### 3 Results

This study assessed a total of 309 pregnant women, all of whom were in the third trimester of pregnancy. Table 1 presents the socio-demographic and obstetrics characteristics of the participants. The mean age of the participants was  $30.49 \pm 5.03$  years. A majority (63.8%) were multigravida, while 27.8% were primigravida. Most participants (62.8%) initiated antenatal care during the second trimester. The overwhelming majority (91.6%) were married, and 91.5% identified as Yoruba. Among the participants, 77.0% had

**Table 1.** Participants’ Demographic and Obstetric Characteristics

Variable	Frequency	Pourcentage(%)	Mean ± SD
<b>Age (years)</b>			30.49 ± 5.03
<b>Age group (years)</b>			
- Less than 24	32	10.4	
- 25-29	111	35.9	
- 30-34	92	29.8	
- 35 and above	74	23.9	
<b>Marital status</b>			
- Single or Serated	26	8.4	
- Married	283	91.6	
<b>Religion</b>			
- Christianity	176	57.0	
- Income < Expenses	60 (37)	20 (38.5)	
<b>Level of education</b>			
- None and primary	6	1.9	
- secondary	65	21.0	
- Tertiary	238	77.0	

**Table 1.** Continued

Variable	Frequency	Pourcentage(%)	Mean ± SD
<b>Ethnicity</b>			
- Yoruba	283	91.6	
- Others	26	8.4	
<b>Family Income</b>			
- None	19	6.1	
- Mow income	88	28.5	
- Middle income	166	53.7	
- Hign income	36	11.7	
<b>Gravidity</b>			2.46 ± 1.23
- Primigravida	86	27.8	
- Multi-gravida	197	63.8	
- Grand multi-gravida	26	8.4	
<b>Parity</b>			1.05 ± 1.01
- Nulliparity	131	42.4	
- Multiparity	178	57.6	
<b>GA at booking</b>			
- 1 <sup>st</sup> Trimester	83	26.9	
- 2 <sup>nd</sup> Trimester	194	62.8	
- 3 <sup>rd</sup> Trimester	32	10.4	
<b>Study center (Health facility)</b>			
- Tertiary	161	52.1	
- Secondary	148	47.9	

Note. \*GA – Gestational age

attained tertiary education, 53.7% fell within the middle-income category, and 57.0% were Christians. The mean gestational age at enrolment was 33.4 weeks (SD= 3.60).

### 3.1 Micronutrient supplements intake among the pregnant women

There intake of most of the micronutrient supplements among the pregnant women was suboptimal. Approximately half of the participants (51.1%) demonstrated adequate intake of ferrous supplements, while 44.7% had sufficient intake of vitamin C (Table 2). Although a significant majority (73.5%) achieved adequate intake of folic acid, only 46 % reported adequate consumption of vitamin B-complex. In contrast, fewer than 30% of the participants exhibited adequate intake of calcium (23.0%), multivitamin (15.2%),

**Table 2.** Proportion of participants who had adequate weekly micronutrient supplement intake

Weekly Intake	Number of participants with adequate intake	Percentage
Folic acid	227	73.5
Vitamin B-complex	142	46.0
Ferrous	158	51.1
Vitamin C	138	44.7
Multivitamin	47	15.2
Omega-3	36	11.7
Vitamin A	14	4.5
Zink	28	9.1
Calcium	71	23.0
Vitamin D	21	6.8

Note. Adequate intake implies 4 or more consumptions per week

omega-3 (11.7%), zinc (9.1%), vitamin D (6.8%) and vitamin A (4.5%). (Table 2).

### 3.2 Change in PCV levels after receiving ANC care

Figure 1 illustrates the changes in hematocrit levels among participants, comparing the prevalence of anemia at the initial antenatal care (ANC) booking to that observed at the time of enrolment in the study after receiving ANC care, including hematinic supplementation. The average duration of supplement intake was 14.9 weeks (SD=6.32). At booking, 42.9% of participants were classified as anemic, while this proportion significantly declined to 26.8% by the time of enrolment in the third trimester. Table 3 highlights the mean PCV levels at enrolment (Mean=34.03, SD=2.90) compared to booking (Mean=33.46, SD=3.83); with the difference being statistically significant ( $p = 0.038$ ). This indicates a notable


**Table 3.** Test of change in PCV level using paired samples *t*-test

Repeated Measures	Mean	SD	<i>t</i>	Df	Effect size	<i>p</i> -value
PCV level at enrolment	34.03	2.903	2.089	155	155	0.038**
PCV level at booking	33.46	3.826				

Note. \*\* Significant at 5% level of significance ( $p < 0.05$ ); Df: Degree of Freedom

improvement in PCV levels following ANC care. Moreover,

91% of women classified as anemic in the third trimester had mild anemia, with only 4% experiencing severe anemia.



**Figure 1.** Comparing prevalence of anemia at point of booking and enrolment (Interview)

### 3.3 Association between demographic and obstetrics characteristics and changes in PCV levels

Linear regression analysis revealed significant associations between participants' demographic and obstetric characteristics and changes in PCV levels. Age group 25 – 29 years ( $\beta = 1.186$ ,  $p$ -value = 0.029, CI of  $\beta = 0.124 - 2.247$ ), level of education ( $\beta = 2.734$ ,  $p$ -value = 0.041, CI of  $\beta = 0.118 - 5.349$ ) and religion ( $\beta = -0.918$ ,  $p$ -value = 0.007, CI of  $\beta = -1.586 - -0.251$ ) were significantly associated with increases in PCV levels. However, factors such as income level, gravidity, parity, use of intermittent preventive therapy for malaria (IPTp), and gestational age at booking did not show significant association with changes in hematocrit levels.

Specifically, participants aged 25 – 29 years experienced a 1.19-unit greater increase in PCV levels in the third trimester (at enrolment) after receiving ANC care, including hematinic supplementation, compared to those under 25 years. However, the change in PCV levels among participants aged 30 – 34 years and 35 years or older were not significantly different from those observed in the group below 25 years. Furthermore, participants with a tertiary education level showed a 2.73-unit greater increase in PCV levels after ANC care compared to those with primary education or no formal education. Meanwhile, participants practicing Islam demonstrated a lower effect (-0.918) of PCV level improvement following ANC compared to their Christian counterparts (Table 4).

### 3.4 Association between micronutrient supplement intake and changes in PCV Levels after a period of ANC care

A model diagnostic test was performed to evaluate the fitness of the Linear Mixed Model (LMM). The diagnostic process included examining a histogram of error terms and performing the Shapiro-Wilk test. The histogram indicated approximately normally distributed error terms, a finding supported by the Shapiro-Wilk test ( $statistic = 0.935$ ,  $p = 0.143$ ). Similarly, the fitness of the Linear Regression Model for Repeated Measures (LRMRM) was assessed using the same diagnostic methods. The histogram again suggested approximately normal distribution of error terms, corroborated by the Shapiro-Wilk test ( $statistic = 0.993$ ,  $p = 0.215$ ).

The analysis revealed that adequate intake of folic acid, vitamin B-complex, vitamin C, omega-3, zinc, and calcium appeared to improve the PCV levels of participants. Conversely, the intake of ferrous supplements, multivitamins, vitamin A, and vitamin D did not demonstrate a positive effect. However, none of these associations reached statistical significance, as indicated in Table 5.

**Table 4:** Demographic and Obstetrics Factors Influencing Change in PCV Level Using Linear Mixed Model (LMM)

Control Factor	Estimate	Std. Err	z-value	Partial Eta Squared	p-value	95% Confidence Interval	
						Lower Limit	Upper Limit
<b>Study Centre</b>							
- Centre 1 = UCH	0.684	0.317	2.157	<b>0.015</b>	<b>0.032*</b>	0.061	1.307
- Centre 2 = Adeoyo	1.03	0.05	20.07	0.00	0.89	0.80	
<b>Fixed Factor</b>							
<b>Age (years)</b>							
- 35 and above	0.834	1.592	1.324	0.006	0.186	-0.404	2.072
- 30-34	0.478	0.630	0.841	0.002	0.401	-0.639	1.595
- 25-29	1.186	0.568	2.195	<b>0.016</b>	<b>0.029*</b>	0.124	2.247
- Less than 25	0						

Table 4: Continued

Control Factor	Estimate	Std. Err	z-value	Partial Eta Squared	p-value	95% Confidence Interval	
						Lower Limit	Upper Limit
<b>Marita Status</b>							
- Married	0.390	0.560	0.697	0.002	0.486	-0.710	1.491
- Single							
<b>Level of Education</b>							
- Tertiary	2.734	1.331	2.054	<b>0.012</b>	<b>0.041*</b>	0.118	5.349
- Secondary	1.818	1.370	1.327	0.006	0.185	-0.875	4.511
- Primary or none	0						
<b>Religion</b>							
- Islam	-0.918	0.340	-2.705	<b>0.025</b>	<b>0.007**</b>	-1.586	-0.251
- Christianity	0						
<b>Ethnicity</b>							
- Others	-0.9069	0.585	1.658	0.009	0.098	-0.180	2.119
- Yoruba	0						
<b>Gravidity at Booking</b>							
- Grand multigravida	0.949	0.759	1.250	???	0.212	-0.543	2.441
- Multi-gravid	0.880	0.505	1.742	0.005	0.082	-0.113	1.873
- Primigravid	0			0.010			
<b>Parity</b>							
- Multipara	-0.784	0.470	-1.668	0.009	0.096	-1.708	0.140
- Nullipara	0						
<b>GA at Booking</b>							
- 3 <sup>rd</sup> Trimester	-0.238	0.595	-0.399	0.001	0.690	-1.408	0.999
- 2 <sup>nd</sup> trimester	0.212	0.362	0.585	0.002	0.559	-0.500	0.924
- 1 <sup>st</sup> Trimester	0						

Note. \* Significant at 5% level of significance; \*\* significant at 1.7% level of significance (Bonferroni Adjusted *p*-value)

**Table 5:** Association between micronutrient supplement intake and change in PCV level using linear regression model for repeated measures (LRMRM)

Micronutrient Intake	Estimate	Std. Err	z-value	Partial Eta Squared	p-value	95% Confidence Interval	
						Lower Limit	Upper Limit
<b>Folic Acid</b>							
- Adequate	0.106	0.456	0.233	0.002	0.816	-0.791	1.003
- Poor	0						
<b>Vitamin B-complex</b>							
- Adequate	0.104	0.415	0.250	0.002	0.803	-0.712	0.919
- Poor	0						
<b>Ferrous</b>							
- Adequate	-0.808	0.400	-0.771	0.0019	0.441	-1.095	0.478
- Poor	0						
<b>Vitamin C</b>							
- Adequate	0.148	0.391	0.377	0.0005	0.706	-0.622	0.917
- Poor	0						
<b>Multivitamin</b>							
- Adequate	-0.180	0.538	-0.334	0.0004	0.738	-1.237	0.879
- Poor	0						
<b>Omega3</b>							
- Adequate	0.599	0.606	0.988	0.0031	0.324	-0.593	1.791
- Poor	0						

Table 5: Continued

Micronutrient Intake	Estimate	Std. Err	z-value	Partial Eta Squared	p-value	95% Confidence Interval	
						Lower Limit	Upper Limit
<b>Vitamin A</b>							
- Adequate	-0.560	1.098	-0.510	0.0008	0.610	-2.719	1.599
- Poor	0						
<b>Zinc</b>							
- Adequate	0.482	0.706	0.681	0.0016	0.496	-0.909	1.873
- Poor	0						
<b>Calcium</b>							
- Adequate	0.574	0.478	1.200	0.0048	0.231	-0.366	1.514
- Poor	0						
<b>Vitamin D</b>							
- Adequate	-0.849	0.843	-1.007	0.0035	0.314	-2.506	0.808
- Poor	0						

Note. \* Significant at 5% level of significance; \*\* Significant at 1.7% level of significance (Bonferroni Adjusted p-value)

### 3.5 Association between intake of diets rich in agents that impair iron absorption and change in PCV levels

Table 6 depicts the association between the intake of diet rich in agents that impair iron absorption and changes in PCV

Table 6: Association between Intake of diet rich in agents that impair Fe absorption and change in PCV levels

Micronutrient Intake	Estimate	Std. Err	z-value	Partial Eta Squared	p-value	95% Confidence Interval	
						Lower Limit	Upper Limit
<b>Milk</b>							
- Adequate	-0.719	0.360	-1.999	<b>0.0138</b>	<b>0.046*</b>	-1.426	-0.012
- Poor	0						
<b>Soya beans</b>							
- Adequate	-0.334	0.603	-0.554	0.0013	0.580	-1.520	0.852
- Poor	0						
<b>Tea</b>							
- Adequate	0.351	0.413	0.850	0.0028	0.396	0.461	1.163
- Poor	0						
<b>Chocolate</b>							
- Adequate	-0.304	0.359	-0.847	0.0025	0.397	-1.009	0.401
- Poor	0						

Note. Poor intake implies consumption less than 4 days per week. \* Significant at 5% level of significance \*\* significant at 1.7% level of significance (Bonferroni Adjusted p-value)

levels. The analysis indicated that adequate milk intake ( $\beta = -0.819$ ,  $p = 0.029$ , CI of  $\beta = 0.124 - 2.247$ ) had a significant negative effect on change in PCV level, while adequate intake of soya beans ( $\beta = -0.334$ ,  $p = 0.580$ , CI of  $\beta = -1.520 - 0.852$ ) and cocoa-based beverages ( $\beta = -0.304$ ,  $p = 0.397$ , CI of  $\beta = -1.009 - 0.401$ ) demonstrated a negative effect on changes in hematocrit levels; however, these effects were not statistically

significant. Conversely, tea intake ( $\beta = 0.351$ ,  $p = 0.396$ , CI of  $\beta = -0.461 - 1.163$ ) was associated with a positive effect on hematocrit changes, though this effect also did not reach statistical significance.

## 4 Discussion

This study evaluated the micronutrients supplementation among pregnant women receiving ANC and its impact on hematocrit levels. Key findings include: approximately three-quarters of the participants initiated ANC in the second trimester or later, and around two in five women were anemic

at the time of booking. While, about half of the participants demonstrated adequate intake of iron supplements, three-quarters reported adequate folic acid intake. However, fewer than half had adequate vitamin B-complex intake, and only a minority used additional multivitamins, vitamin A, vitamin D, or calcium supplements. Adequate intake was defined as supplement use for at least four days per week. By the time of the study interview, less than one-third of the participants were anemic, indicating improved hematocrit levels during the third trimester.

#### 4.1 Sociodemographic and obstetric characteristics

The mean age of participants in this study was 30.49 years ( $\pm 5.03$ ), consistent with findings from studies conducted in Enugu and Algeria that reported mean ages of 30.2 ( $\pm 5.2$ ) years and 30.3 ( $\pm 6.2$ ) years, respectively (Bayazid et al., 2021; Dim & Onah, 2007). This is expected, as the participants were pregnant women within the reproductive age group and predominantly multiparous.

Unlike studies in Uyo and Abeokuta, which found associations between obstetric factors (e.g., gestational age, and use of intermittent preventive treatment in pregnancy [IPTp]) and anemia, this study observed no significant association between these factors and low hematocrit levels (Idowu et al., 2005; Olatunbosun et al., 2014). This discrepancy could be attributed to the higher socioeconomic status of the study participants, which likely ensured food security and access to balanced diets rich in essential micronutrients.

#### 4.2 Prevalence of Anemia and Hematocrit levels

At booking, the prevalence of anemia was 42.9%, higher than the 32.5% reported in Sagamu by Sholeye et al. (2017) but lower than the 54.5% reported in Uyo by Olatunbosun et al., (2014). Over 90% of anemic participants in this study had mild anemia, exceeding the 70% reported in Sagamu (Olatunbosun et al., 2014; Sholeye et al., 2017). Few participants experienced moderate to severe anemia. These variations may be due to differences in study settings; this study included secondary and tertiary healthcare facilities with participants of diverse socioeconomic backgrounds, whereas the Sagamu study was conducted in primary healthcare centers, where low socioeconomic status and food insecurity were prevalent.

Age and educational status were significantly associated with hematocrit levels in the current study, aligning with findings from previous studies (Olatunbosun et al., 2014; Omote et al., 2020). Religion was also significantly associated with hematocrit levels, though this study could not establish a clear

association between Islamic practice and anemia. Sociocultural and religious dietary restrictions may influence nutritional intake and anemia, as documented in previous studies (Major-Smith et al., 2023). For instance, vegetarians avoid animal protein sources rich in iron and, and members of certain religious groups, such as Jehovah's Witnesses, abstain from blood transfusions even in severe anemia, instead prioritizing hematinic supplementation (Major-Smith et al., 2023; Berg et al., 2022). Interestingly, adequate micronutrient supplementation alone was not significantly associated with positive changes in PCV levels. This finding aligns with earlier studies suggesting that socioeconomic status and food security are critical determinants of anemia in pregnancy (Olatunbosun et al., 2014; Omote et al., 2020). Some women may compensate for inconsistent supplementation with optimal dietary intake, while others may rely heavily on supplements due to dietary inadequacies. The multifactorial nature of anemia is evident in this study. Factors such as the socioeconomic status (closely associated with education level), healthcare facilities, age, and possibly preconception use of hematinics, as well as booking gestational age, may have contributed to the observed outcomes.

#### 4.3 Micronutrient supplementation and dietary intake

The micronutrient supplements most commonly consumed by the participants included folic acid, iron, vitamin C and B-complex tablets. These supplements constitute the primary hematinics routinely prescribed to pregnant women by antenatal healthcare providers. However, the frequency of micronutrients intake among participants was suboptimal, consistent with findings from Bayazid et al., (2021) in western Algeria, but lower than the rates reported by Amadi-Joy et al., (2017) in Imo, Nigeria, where over 80% of participants adhered to daily intake of iron, folic acid, vitamin C, and B-complex tablets. Similarly, Enyew et al. (2023) observed suboptimal micronutrient intake among pregnant women in East Africa. However, evidence indicates that optimal prophylactic iron supplementation can reduce maternal anemia at term by 70% (Peña-Rosas et al., 2015).

The intake of other critical micronutrients, such as calcium, vitamins A and D, and omega 3 fatty acids, was also low, consistent with findings by Amadi-Joy et al., (2017). According to the WHO recommendations, these supplements especially calcium, zinc, vitamin A and Vitamin D should be incorporated into routine antenatal care in regions with documented deficiencies (Amadi-Joy et al., 2017; WHO, 2020). Despite evidence of high prevalence of deficiencies across different socioeconomic groups in Nigeria, the utilization of these micronutrients remains insufficient. For instance, calcium deficiency prevalence ranges from 29.2



to 58.76%, vitamin A deficiency from 21 to 35%, and vitamin D deficiency is from 4.8 to 32.3% (Ajong *et al.*, 2019; Bako *et al.*, 2021; Hanson *et al.*, 2019; Owie & Afolabi, 2018).

Adequate supplementation with these micronutrients is necessary, not only to prevent deficiencies but also to prevent potential fetomaternal complications. Deficiencies in these nutrients have been associated with congenital malformations, cretinism, preterm birth, low birthweight, intrauterine growth restriction, pre-eclampsia, and inadequate lactation that may arise from their deficiencies (Gernand *et al.*, 2016; Harika *et al.*, 2017; Parisi *et al.*, 2019).

The ingestion of substances such as milk, cocoa or soy products, which are known to impair iron absorption, was associated with limited improvements in hematocrit levels. Among these, only adequate milk intake was significantly associated with negative hematocrit changes, reinforcing the inhibitory role of certain compounds in milk on iron absorption (Al Hasan *et al.*, 2016; Armah *et al.*, 2015). Studies have shown that milk, cocoa, tea or soya beans contain either phytates, polyphenol, oxalic acid which chelates iron and impair its absorption. Nevertheless, as suggested by some authors, effects of intake of these iron inhibitors on availability of iron for hemopoiesis in the body is dose dependent and are countered by effect of iron absorption enhancing substances such as ascorbic acid and animal source food which are also contained in the diets (Al Hasan *et al.*, 2016; Armah *et al.*, 2015; Björn-Rasmussen & Hallberg, 1979; He *et al.*, 2018).

#### 4.4 Strategies for addressing anemia in pregnancy

To mitigate anemia during pregnancy, several strategies should be implemented. First, comprehensive health education programs targeting young women are essential to emphasize the importance of early antenatal booking and the timely initiation of hematinic supplements. Specific health talks should be designed to promote the intake of iron, folic acid, vitamins A, B12, and C as well as multiple micronutrient supplement (MMS) as recommended by the WHO.

Early detection and prompt management of anemia at the initial antenatal visit can significantly improve hematocrit levels, reducing the prevalence of anemia by the time of delivery. Additionally, the implementation of effective malaria preventive measures, including the use of insecticide-treated nets (ITNs) and IPTp, will also reduce incidence of anemia caused by malaria in pregnancy. The prophylactic treatment of helminthiasis in areas with high prevalence can reduce anemia caused by parasitic infections.

Counselling women on family planning to encourage adequate inter-pregnancy intervals using modern contraceptives and uptake of other components of preconception care can provide women with sufficient time to restore optimal hematocrit levels before conception.

Socioeconomic factors also play a critical role. Policies aimed at providing employment opportunities for women, subsidizing food prizes to enhance access to nutritious meals, and promoting food fortification with essential micronutrients are vital to improving hematocrit levels during pregnancy women and preventing adverse obstetric outcomes related to anemia (Cappellini & Motta, 2015; Jugha *et al.*, 2020; Lebso *et al.*, 2017; Ramachandran, 2021; WHO, 2020; Zerfu *et al.*, 2016).

#### Limitations of the study

This study is not without some limitations. Most participants initiated antenatal care late, typically in the second trimester, and nearly half were anemic at the time of booking. A longitudinal study would have provided more comprehensive insights into the progression of hematocrit levels across trimesters of pregnancy. Such an approach would have also allowed for a more robust analysis of the relationship between the dose and duration of micronutrient supplementation and changes in hematocrit levels. The study was unable to ascertain the proportion of women who had initiated micronutrient supplements prior their antenatal care booking. Additionally, reliance on participants' self-reported dietary habits and compliance with micronutrient supplementation introduces potential recall bias. Conducting a complete blood count for all participants, complemented by peripheral blood smear analysis for those with low hematocrit levels, could have provided valuable information regarding the type, cause and severity of anemia.

Despite its limitations, this study offers valuable insights into the patterns of micronutrient supplementation among pregnant women, including less commonly discussed supplements such as zinc and vitamins A and D, and their impact on hematocrit levels.

## 5 Conclusions

In conclusion, despite the routine prescription of micronutrient supplements by antenatal care providers, the participants demonstrated suboptimal intake of essential hematinics, including iron and vitamin B complex, as well as other key supplements such as vitamins A, D, calcium, multivitamins, omega-3, and zinc. The prevalence of anemia in the third trimester was 26.8%, remains a significant concern for antenatal and delivery healthcare providers.

The findings indicate that low hematocrit levels during pregnancy are associated with demographic and

socioeconomic factors, including maternal age, education level, and food security. It is imperative that every antenatal care visit serves as an opportunity to enhance patient education and counselling regarding the benefits of maintaining optimal hematocrit levels and the risk associated with anemia during pregnancy.

The frequency of intake and compliance with intake of hematinic and micronutrients must be assessed at every antenatal clinic visit and impeding factors must be addressed to improve the utilization of these micronutrient supplements among pregnant women. While self-reported adherence is subjective, introducing frequency logs for micronutrient intake could serve as a motivational and monitoring tool, encouraging compliance and facilitating more effective interventions.

**Acknowledgment:** None.

**Source of funding:** This study has not been funded by any institution.

**Previous submissions:** This work has not been published previously in any journal or in any other form.

**Authors' Contribution:** **Abdus-Salam:** Conceptualization, Research development, data analysis, manuscript writing. **Oni:** Conceptualization, Research development, data analysis, manuscript writing. **Adeyanju:** Conceptualization, Research development, data analysis, manuscript writing. **Adeyeye, Victoria, Oluwatosin, Adebayo-Tayo, Medebem, Tongo, Ajagbe, Nwaigwe, Akinosi, and Oladayo:** Conceptualization, Research development, data collection.

\***College of Medicine, University of Ibadan, Ibadan Medical Students Group B 2019:** Adeyeye Oluwabunmi, Ajani Victoria A, Taofeeq Oluwatosin T, Adebayo-Tayo Mercy A, Medebem Obiageli O, Tongo Esosa S, Ajagbe Adenike O, Nwaigwe Amarachi P, Akinosi Taiwo M, Olatunbosun Oladayo D, Aba Oluwafemi E, Adebisi Abeeb A, Adedeji Praise K, Adesioye Andrew O, Afigbo Wisdom C, Anyam Ngusha V, Apologun Upe E, Arogundade Zainab B, Audu Victor, Awokunle Joshua A, Babalola Adedoyin V, Ezech Henry C, Fashina Oluwadarasimi T, Folarin Joshua O, Folorunsho Ajibola A, Jonah Caleb O, Kifodu Ugochukwu J, Lateef Bisola M, Nebo Ifunanyachukwu T, Nkeakam Fortune A, Nwamadigesi Caroline A, Obitayo Sunday O, Odunsi Temidayo, Ogunnaike Gbemisola B, Ojuro Vincent C, Oladepo Olayinka, Olorunfemi Victor, Onuoha Obianuju, Onyishi Martin I and Oyinola Priscilla J.

**Conflicts of Interest:** All authors declare that they have no conflicts of interest.

**Ethical approval:** This study was approved by the institutional ethics Review Committee of University College Hospital and the College of Medicine, University of Ibadan (UI/UCH Ethics Committee–UI/EC/23/0040). Voluntary and informed consent of the participants was obtained.

**Statement of Informed Consent:** All participants in this study gave voluntary informed consent.

**Preprint deposit:** Authors did not share this manuscript as a preprint deposit.

## References

- Abdallah, F., John, S. E., Hancy, A., Paulo, H. A., Sanga, A., Noor, R., Lankoande, F., Chimanya, K., Masumo, R. M., & Leyna, G. H. (2022). Prevalence and factors associated with anaemia among pregnant women attending reproductive and child health clinics in Mbeya region, Tanzania. *PLOS Global Public Health*, 2(10), e0000280. <https://doi.org/10.1371/journal.pgph.0000280> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Agbozo, F., Abubakari, A., Der, J., & Jahn, A. (2020). Maternal Dietary Intakes, Red Blood Cell Indices and Risk for Anemia in the First, Second and Third Trimesters of Pregnancy and at Predelivery. *Nutrients*, 12(3), 777. <https://doi.org/10.3390/nu12030777> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Ajong, A. B., Kenfack, B., Ali, I. M., Yakum, M. N., & Telefo, P. B. (2019). Prevalence and correlates of low serum calcium in late pregnancy: A cross sectional study in the Nkongsamba Regional Hospital; Littoral Region of Cameroon. *PLoS ONE*, 14(11), e0224855. <https://doi.org/10.1371/journal.pone.0224855> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Al Hasan, S. M., Hassan, M., Saha, S., Islam, M., Billah, M., & Islam, S. (2016). Dietary phytate intake inhibits the bioavailability of iron and calcium in the diets of pregnant women in rural Bangladesh: A cross-sectional study. *BMC Nutrition*, 2(1), 24. <https://doi.org/10.1186/s40795-016-0064-8> [Crossref] [Google Scholar] [Publisher]
- Amadi-Joy, A., Asinobi, C., & Afem-Anene, O. (2017). Intake of multiple micronutrients supplementation during pregnancy and effects on birth weight of infants in Umuna, Orlu Local Government Area, Imo State, Nigeria. *Journal of Dietitians Association of Nigeria*, 8, 25–33. [Google Scholar] [Publisher]
- Armah, S. M., Boy, E., Chen, D., Candal, P., & Reddy, M. B. (2015). Regular Consumption of a High-Phytate Diet Reduces the Inhibitory Effect of Phytate on Nonheme-Iron Absorption in Women with Suboptimal Iron Stores. *The Journal of Nutrition*, 145(8), 1735–1739. <https://doi.org/10.3945/jn.114.209957> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Bako, B., El-Nafaty, A. U., Mshelia, D. S., Gali, R. M., Isa, B., & Dungus, M. M. (2021). Prevalence and risk factors of hypocalcemia among pregnant and non-pregnant women in Maiduguri, Nigeria: A cross-sectional study. *Nigerian Journal of Clinical Practice*, 24(1), 75–80. [https://doi.org/10.4103/njcp.njcp\\_640\\_19](https://doi.org/10.4103/njcp.njcp_640_19) [Crossref] [PubMed] [Google Scholar] [Publisher]
- Barrett, J. F., Whittaker, P. G., Williams, J. G., & Lind, T. (1994). Absorption of non-haem iron from food during normal pregnancy. *BMJ (Clinical Research Ed.)*, 309(6947), 79–82. <https://doi.org/10.1136/bmj.309.6947.79> [Crossref] [PubMed] [Google Scholar] [Publisher]

- Bayazid, A., Soum, M., Boumaza, O., & Toumi, H. (2021). Micronutrient supplementation among pregnant women in western Algeria. *The North African Journal of Food and Nutrition Research*, 5(11), <https://doi.org/10.51745/najfnr.5.11.15-22> [Crossref] [Google Scholar] [Publisher]
- Berg, L., Dave, A., Fernandez, N., Brooks, B., Madgwick, K., Govind, A., & Yoong, W. (2022). Women who decline blood during labour: Review of findings and lessons learnt from 52 years of Confidential Enquiries into maternal mortality in the United Kingdom (1962-2019). *European Journal of Obstetrics, Gynecology, and Reproductive Biology*, 271, 20–26. <https://doi.org/10.1016/j.ejogrb.2022.01.028> [Crossref] [Google Scholar] [Publisher]
- Björn-Rasmussen, E., & Hallberg, L. (1979). Effect of animal proteins on the absorption of food iron in man. *Nutrition and Metabolism*, 23(3), 192–202. <https://doi.org/10.1159/000176256> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Cappellini, M. D., & Motta, I. (2015). Anemia in Clinical Practice-Definition and Classification: Does Hemoglobin Change With Aging? *Seminars in Hematology*, 52(4), 261–269. <https://doi.org/10.1053/j.seminhematol.2015.07.006> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Chowdhury H A. (2015). Factors associated with maternal anaemia among pregnant women in Dhaka city, *BMC Women's Health*. <https://bmcwomenshealth.biomedcentral.com/articles/10.1186/s12905-015-0234-x> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Darnton-Hill, I. (2012). Global burden and significance of multiple micronutrient deficiencies in pregnancy. *Nestle Nutrition Institute Workshop Series*, 70, 49–60. <https://doi.org/10.1159/000337421> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Dim, C. C., & Onah, H. E. (2007). The Prevalence of Anemia Among Pregnant Women at Booking in Enugu, South Eastern Nigeria. *Lancet*, 9(3), 11. [PubMed] [Google Scholar] [Publisher]
- Enyew, E. B., Tareke, A. A., Dubale, A. T., Fetene, S. M., Ahmed, M. H., Feyisa, M. S., & Ngusie, H. S. (2023). Micronutrient intake and associated factors among pregnant women in East Africa: Multilevel logistic regression analysis. *PLoS One*, 18(4), e0281427. <https://doi.org/10.1371/journal.pone.0281427> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Frayne, B., Crush, J., & McLachlan, M. (2014). Urbanization, nutrition and development in Southern African cities. *Food Security*, 1(6), 101–112. <https://doi.org/10.1007/s12571-013-0325-1> [Crossref] [Google Scholar] [Publisher]
- Galloway, R. (2003). Anemia prevention and control: What works; part 1: Program guidance. (1–1). USAID, World Bank, UNICEF, PAHO, FAO, MI.
- Gernand, A. D., Schulze, K. J., Stewart, C. P., West, K. P., & Christian, P. (2016). Micronutrient deficiencies in pregnancy worldwide: Health effects and prevention. *Nature Reviews. Endocrinology*, 12(5), 274–289. <https://doi.org/10.1038/nrendo.2016.37> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Hanson, C., Lyden, E., Anderson-Berry, A., Kocmich, N., Rezac, A., Delair, S., Furtado, J., Van Ormer, M., Izevbogie, N., Olateju, E., Akaba, G. O., Anigilaje, E., Yunusa, T., & Obaro, S. (2019). Erratum: Status of Retinoids and Carotenoids and Associations with Clinical Outcomes in Maternal-Infant Pairs in Nigeria. *Nutrients* 2018, 10, 1286. *Nutrients*, 11(2), 396. <https://doi.org/10.3390/nu11020396> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Harika, R., Faber, M., Samuel, F., Kimiywe, J., Mulugeta, A., & Eilander, A. (2017). Micronutrient Status and Dietary Intake of Iron, Vitamin A, Iodine, Folate and Zinc in Women of Reproductive Age and Pregnant Women in Ethiopia, Kenya, Nigeria and South Africa: A Systematic Review of Data from 2005 to 2015. *Nutrients*, 9(10), 1096. <https://doi.org/10.3390/nu9101096> [Crossref] [PubMed] [Google Scholar] [Publisher]
- He, W., Li, X., Ding, K., Li, Y., & Li, W. (2018). Ascorbic Acid can Reverse the Inhibition of Phytic Acid, Sodium Oxalate and Sodium Silicate on Iron Absorption in Caco-2 cells. *International Journal for Vitamin and Nutrition Research. Internationale Zeitschrift Fur Vitamin- Und Ernährungsforschung. Journal International De Vitaminologie Et De Nutrition*, 88(1–2), 65–72. <https://doi.org/10.1024/0300-9831/a000503> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Idowu, O. A., Mafiana, C. F., & Dapo, S. (2005). Anaemia in pregnancy: A survey of pregnant women in Abeokuta, Nigeria. *African Health Sciences*, 5(4), 295–299. <https://doi.org/10.5555/afhs.2005.5.4.295> [PubMed] [Google Scholar] [Publisher]
- Jugha, V. T., Anchang-Kimbi, J. K., Anchang, J. A., Mbeng, K. A., & Kimbi, H. K. (2021). Dietary Diversity and Its Contribution in the Etiology of Maternal Anemia in Conflict Hit Mount Cameroon Area: A Cross-Sectional Study. *Frontiers in Nutrition*, 7, 625178. <https://doi.org/10.3389/fnut.2020.625178> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Kidanto, H. L., Mogren, I., Lindmark, G., Massawe, S., & Nystrom, L. (2009). Risks for preterm delivery and low birth weight are independently increased by severity of maternal anaemia. *South African Medical Journal = Suid-Afrikaanse Tydskrif Vir Geneeskunde*, 99(2), 98–102. [PubMed] [Google Scholar] [Publisher]
- Kish, L. (1965) Survey Sampling. John Wiley and Sons Inc., New York. [Publisher]
- Lebso, M., Anato, A., & Loha, E. (2017). Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PLOS ONE*, 12(12),

- <https://doi.org/10.1371/journal.pone.0188783> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Major-Smith, D., Morgan, J., Emmett, P., Golding, J., & Northstone, K. (2023). Associations between religious/spiritual beliefs and behaviours and dietary patterns: analysis of the parental generation in a prospective cohort study (ALSPAC) in Southwest England. *Public health nutrition*, 26(12), 2895–2911. <https://doi.org/10.1017/S1368980023001866> [Crossref] [PubMed] [Google Scholar] [Publisher]
- National Population Commission (NPC) [Nigeria] and ICF International. (2014). *Nigeria Demographic and Health Survey 2013*. [Google Scholar] [Publisher]
- Olatunbosun, O. A., Abasiattai, A. M., Bassey, E. A., James, R. S., Ibanga, G., & Morgan, A. (2014). Prevalence of Anaemia among Pregnant Women at Booking in the University of Uyo Teaching Hospital, Uyo, Nigeria. *BioMed Research International*, 2014, 849080. <https://doi.org/10.1155/2014/849080> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Omote, V., Ukwamedua, H. A., Bini, N., Kashibu, E., Ubandoma, J. R., & Ranyang, A. (2020). Prevalence, Severity, and Correlates of Anaemia in Pregnancy among Antenatal Attendees in Warri, South-Southern Nigeria: A Cross-Sectional and Hospital-Based Study. *Anemia*, 2020, 1915231. <https://doi.org/10.1155/2020/1915231> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Owie, E., & Afolabi, B. B. (2018). Vitamin D deficiency in pregnant women and newborns in Lagos, Nigeria. *Journal of Obstetrics and Gynaecology: The Journal of the Institute of Obstetrics and Gynaecology*, 38(5), 616–621. <https://doi.org/10.1080/01443615.2017.1396299> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Parisi, F., di Bartolo, I., Savasi, V. M., & Cetin, I. (2019). Micronutrient supplementation in pregnancy: Who, what and how much? *Obstetric Medicine*, 12(1), 5–13. <https://doi.org/10.1177/1753495X18769213> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Peña-Rosas, J. P., De-Regil, L. M., Garcia-Casal, M. N., & Dowswell, T. (2015). Daily oral iron supplementation during pregnancy. *The Cochrane Database of Systematic Reviews*, 2015(7), CD004736. <https://doi.org/10.1002/14651858.CD004736.pub5> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Ramachandran, P. (2021). Prevention & management of anaemia in pregnancy: Multi-pronged integrated interventions may pay rich dividends. *The Indian Journal of Medical Research*, 154(1), 12–15. [https://doi.org/10.4103/ijmr.IJMR\\_994\\_20](https://doi.org/10.4103/ijmr.IJMR_994_20) [Crossref] [PubMed] [Google Scholar] [Publisher]
- Ruiz-Cabello, P., Soriano-Maldonado, A., Delgado-Fernandez, M., Alvarez-Gallardo, I. C., Segura-Jimenez, V., Estevez-Lopez, F., Camiletti-Moirón, D., & Aparicio, V. A. (2017). Association of Dietary Habits with Psychosocial Outcomes in Women with Fibromyalgia: The al-Ándalus Project. *Journal of the Academy of Nutrition and Dietetics*, 117(3), 422–432.e1. <https://doi.org/10.1016/j.jand.2016.09.023> [Crossref] [PubMed] [Google Scholar] [Publisher]
- Sholeye, O. O., Animasahun, V. J., & Shorunmu, T. O. (2017). Anemia in pregnancy and its associated factors among primary care clients in Sagamu, Southwest, Nigeria: A facility-based study. *Journal of Family Medicine and Primary Care*, 6(2), 323–329. [https://doi.org/10.4103/jfmprc.jfmprc\\_74\\_16](https://doi.org/10.4103/jfmprc.jfmprc_74_16) [Crossref] [PubMed] [Google Scholar] [Publisher]
- World Health Organization (WHO). (2011). *Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity*. [Google Scholar] [Publisher]
- World Health Organization (WHO). (2020). *WHO antenatal care recommendations for a positive pregnancy experience: Nutritional interventions update: Multiple micronutrient supplements during pregnancy*. World Health Organization. <http://www.ncbi.nlm.nih.gov/books/NBK560384/> [PubMed] [Publisher]
- World Health Organization (WHO). (2024). *Antenatal iron supplementation. Nutrition Landscape Information System (NLIS). Nutrition and Nutrition-Related Health and Development Data*. [Publisher]
- Zerfu, T. A., Umeta, M., & Baye, K. (2016). Dietary diversity during pregnancy is associated with reduced risk of maternal anemia, preterm delivery, and low birth weight in a prospective cohort study in rural Ethiopia. *The American Journal of Clinical Nutrition*, 103(6), 1482–1488. <https://doi.org/10.3945/ajcn.115.116798> [Crossref] [PubMed] [Google Scholar] [Publisher]